

TABLE OF CONTENTS

	Page
ABSTRACT (THAI).....	I
ABSTRACT (ENGLISH).....	II
ACKNOWLEDGEMENT.....	III
TABLE OF CONTENTS.....	V
LIST OF TABLES.....	VIII
LIST OF FIGURES.....	IX
CHAPTER	
I INTRODUCTION.....	1
1.1 Background and Significance.....	1
1.2 Research Objectives.....	3
1.3 Scope and limitations of the research.....	3
1.4 Expected outcomes.....	4
II LITERATURE REVIEWS.....	5
2.1 Overview of the literature.....	5
2.1.1 Crystallization technologies.....	5
2.1.2 Direct methodological comparisons.....	16
2.2 Protein crystallization.....	16
2.3 Industrial crystallization of protein.....	19
2.4 Papain and crystallization of papain.....	20
2.4.1 Papain.....	20
2.4.2 Papain crystallization review.....	24
2.5 Conclusion.....	27
III RESEARCH METHODOLOGY.....	29
3.1 Chemicals and reagents.....	29
3.2 Methods and apparatus.....	30

TABLE OF CONTENTS (Continued)

	Page
3.2.1 Preliminary lysozyme study to validate SFO setup and phase mapping tools.....	30
3.2.2 Solubility of papain (gravimetric and RI correlation)...	33
3.2.3 Cooling crystallization of papain	34
3.2.4 Antisolvent crystallization of papain	35
3.2.5 Solvent freeze-out crystallization of papain.....	36
3.3 Papain concentration determination and characterization.....	37
3.3.1 Papain activity assay (BAPNA, UV-vis at 410 nm)	37
3.3.2 Crystal characterizations	40
IV RESULTS AND DISCUSSION.....	41
4.1 Preliminary study of SFO crystallization of lysozyme	41
4.1.1 The solubility study by FBRM and SFO crystallizer testing.....	41
4.2 Results of papain crystallization.....	45
4.2.1 Solubility of papain.....	45
4.2.2 Cooling crystallization for papain	46
4.2.3 Antisolvent crystallization for papain	48
4.2.4 SFO crystallization for papain.....	51
4.2.5 Powder X-ray diffraction	55
4.2.6 SEM of papain crystal.....	56
4.2.7 The enzyme activity analysis.....	57
V CONCLUSION AND RECOMMENDATION.....	61
5.1 Conclusion.....	61
5.2 Recommendations.....	61
REFERENCES	63
APPENDIX A	72
APPENDIX B	74

TABLE OF CONTENTS (Continued)

	Page
BIOGRAPHY	88

LIST OF TABLES

Table		Page
2.1	Summary comparison of crystallization methods.	16
2.2	Papain enzymatic activity studies.	23
4.1	The percentage loss and recovery of papain crystallization using SFO in this work.....	54
4.2	The enzyme activity assay of papain crystal.....	59

LIST OF FIGURES

Figure	Page
1.1 Overall workflow of papain crystallization.	4
2.1 Conceptual solubility and nucleation diagram for cooling crystallization. Adapted from (Giulietti et al., 2001).....	8
2.2 Conceptual solubility/nucleation diagram for the antisolvent technique. Adapted from (Jia et al., 2022).	10
2.3 SEM photomicrographs of nanoparticle papain precipitated at different solvent-to-antisolvent ratios (1:4); reproduced from with permission (Boonkerd et al., 2024; Boonkerd & Wantha, 2024).....	11
2.4 Set up of a simple equipment for solvent freeze out crystallization of proteins. Adapted from (Borbón & Ulrich, 2012).	12
2.5 Mechanism of the solvent freeze out (SFO) crystallization process. Two crystallization processes take place. One is the crystallization of melt (water-ice) on a cooled surface. The other is the crystallization of the protein within the solution. Adapted from (Borbon & Ulrich, 2013).	13
2.6 Effect of salt on macromolecule solubility (Mcpherson, 1990).....	17
2.7 Typical phase diagram with crystallization window where the optimal crystallization region is found near the metastable liquid-liquid equilibrium (LLE) (Curtis & L.Lue, 2005).	18
2.8 Ribbon representation of the three-dimensional structure of papain active site at His159 and Cys25 (Amri & Mamboya, 2012; Grzonka et al., 2007; Kamphuis et al., 1984).....	20
2.9 Commercial enzyme Papain brand "Paya" reaction with casein (Mathias Elsson et al., 2019).....	21

LIST OF FIGURES (Continued)

Figure	Page
2.10 Needle-like crystals of papain present in the suspension originating from Sigma. The needles contain parallel fibrils, which exhibit a pronounced zigzag helical periodicity. Negatively stained with 2% sodium phosphor tungstate (pH 7.0). (a) x18,000, (b) x78,000, (c) x78,000, (d) x6,000, (e) x30,000, respectively. Adapted from (Harris, 1983).....	24
2.11 Crystals of papain protease inhibitor (PPI). (a) Crystal form I (prismatic) grown in 2.0 M sodium formate, 20 mM sodium acetate pH 4.6. (b) Original crystals of crystal form II grown at 293 K in 1.8 M ammonium sulfate, 100 mM MES pH 6.5, 10 mM cobalt chloride. (c) Typical growth crystals of crystal form II of PPI grown at 283 K after optimization. The clusters of needles are only loosely attached to each other and the large single crystal in the middle was picked out for data collection. Adapted from (Mohamed Azarkan et al., 2006).....	25
2.12 Phase-diagrams of papain: Red solubility curve was determined from the residual concentration in equilibrium with crystals 50-days after the initiation of crystallization at varying temperatures from 4 °C, 12 °C and 22 °C, left to right respectively. Nucleation and precipitation curves are plotted in black and green, respectively. Adapted from (Yi-Bin Lin et al., 2008).....	26
2.13 a) XRD pattern of papain crystals under different precipitation conditions. b) SEM micrographs showing crystal morphology of papain under combined precipitant. Adapted from (Qi Hao et al., 2024).....	27
3.1 Papain powder from a) Shaanxi Yuantai Biological Technology Co., Ltd (YT0829), b) Tianjin HEOWNS Biochemical Technology Co., Ltd (P-01102), c) Sisco Research Laboratories Pvt. Ltd (SRL) (papain 2xUSP, 14049) and d) Lysozyme powder from Sigma-Aldrich.	30
3.2 The Particle-Track™ G600B with FBRM (technology from METTLER TOLEDO) and its working principle.....	31

LIST OF FIGURES (Continued)

Figure	Page
3.3	The process of solubility measurement for lysozyme and crystallization by the SFO technique. 32
3.4	SFO crystallizer setup. 33
3.5	Refractive index (RI) vs. papain concentration (g/mL, g/g) in DI water, acetate buffer (pH 5.0), and buffer-methanol mixtures..... 34
3.6	The process of solubility measurement for papain by gravimetric method. At -8, 0, 10, 20, 30 °C, in water, acetate buffer, and buffer mixed methanol (0–60 % w/w). Using a 0.22 µm syringe filter prior to drying. 34
3.7	Cooling crystallizer set up. 35
3.8	Illustration the SFO crystallization procedures for papain..... 37
3.9	UV–vis spectrophotometer principle and equipment. Adapted from (https://microbiologynote.com/spectrophotometer-principle/)..... 38
3.10	The calibration line of papain concentration in water for enzyme activity measurement..... 39
3.11	The papain activity measurement procedure with BAPNA hydrolyze. 39
4.1	The crystallizer with FBRM set-up for measuring lysozyme solubility. 41
4.2	Solubility of lysozyme in ammonium sulfate with various temperatures (a-b). 42
4.3	The phase-diagrams of lysozyme in ammonium sulfate solution. Solubility and nucleation point using SFO at pH 4.6 with salt concentration of (a) 0.4M, (b) 0.6M, and (c) 0.7M. (d) Freeze-coil and bulk-solution temperature profiles with time. 43
4.4	(a) The SFO crystallizer at Lysozyme precipitated as a gel in ammonium sulfate (0.7M) from solution at 1 °C. (b) particle images by Microscope,4X. (c) Filtration and dry particle. 44

LIST OF FIGURES (Continued)

Figure	Page
4.5	The Lysozyme nucleation in ammonium sulfate (0.6M) from SFO crystallization: (a) particle images by Microscope 40X, and (b) Lysozyme cloud in solution but d_1 as a first freezing, d_2 as second freezing. 44
4.6	The solubility of papain in (a) water and acetate buffer (pH 5.0) at various temperatures. (b) In the mixture of acetate buffer with methanol (0.07-0.6 w/w) at various temperatures. 46
4.7	The phase-diagram for cooling crystallization of papain..... 47
4.8	Cooling crystallization (cooling from 30 °C to 12.5 °C) of a saturated papain solution: (a) solution clouding in the crystallizer, (b) photomicrograph of crystals at the onset of nucleation, and (c) photomicrograph of crystals after 24 hrs of nucleation and growth. 48
4.9	Cooling crystallization (cooling from 20 °C to -3 °C) of a saturated papain solution: (a) crystal clouding in the crystallizer after nucleation, (b) photomicrograph of crystals at the onset of nucleation, and (c) photomicrograph of crystals after 24 hrs of nucleation and growth..... 48
4.10	The phase diagram of antisolvent crystallization for papain at 0 °C (a) and 20 °C (b)..... 49
4.11	Antisolvent crystallizer setup with methanol dropping. 49
4.12	(a) Crystallizer image was captured during methanol addition at 0.348 w/w, and photomicrograph of papain crystals in liquor (b) at cloudy point (methanol 0.348 w/w) from saturation concentration of 0.17 g/mL (0 °C); (c) at point of methanol 0.56 w/w, and (d) the particle became sticky gel at 0.57 w/w. 50
4.13	(a) Crystallizer captures image was captured during methanol addition at supersaturation of 0.23 w/w, and photomicrograph of papain crystals in liquor (b) at cloudy point (0.23 w/w) from saturation concentration of 0.67 g/mL (20 °C) and (c) the particle became sticky gel at 0.64 w/w. 50

LIST OF FIGURES (Continued)

Figure	Page
4.14	(a) The phase diagram of SFO for this papain solution at pH 5.0. (b) the temperature reduction profile of freeze coil (293.15 & 311.32h are time of nucleation). *Ft is final temperature of freeze coil. 51
4.15	The SFO crystallizer and photomicrographs of papain crystal obtained at nucleation point from feed concentrations ranging of 0.33 to 0.47 g/mL..... 52
4.16	Examples of photographs of crystals from all conditions obtained after drying on slide under microscope (room/air temperature) in SUT and TJU. 53
4.17	PXRD patterns of papain particle obtained from this work: orange) Amorphous papain from antisolvent by methanol; red) Plate-like form from cooling crystallization; green) Mixed needle/plate like habit crystal of papain from SFO process; yellow) Mix form crystal from cooling crystallization without buffer; purple) Mix form crystal from cooling crystallization with buffer (pH 5.0); and commercial crystalline papain: blue) As product no. P-01102; black) YT0829; dark red) 14049. 55
4.18	The SEM photographic of this work: (a) Commercial crystalline papain, (b-e) Crystal of papain from SFO process. (c) Plate-like form and (d-e) Needle-like form..... 57
4.19	The product from reaction of papain with substrate (BAPNA) was analyzed using UV-vis spectroscopy at 410 nm..... 59
4.20	The papain activity denatured by time in different phases storage inside refrigerator (solid commercial papain and dissolution in water as liquid) and comparison with products of using antisolvent by methanol, SFO process: (a) as BAPNA Units, (b) as Units/mg. The activity denatured in liquid phase: (c) as BAPNA Units, (d) as Units/mg. 60